

for Tomatoes and Vegetables

Sprays in all directions

- even upside down for hard to reach areas -

PROVIDES RAPID CONTROL of Insects Listed on Side Panel MAY BE USED ON FOOD GROPS **TO WITHIN ONE DAY OF HARVEST**

Intended for Residential Use Only





0.2%

EPA REG. NO. 2724-568-72804 EPA EST. NO. 11623-GA-1

100.0%

NET CONTENTS: 16oz. KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional precautionary statements.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & ANIMALS CAUTION

Harmful if absorbed through skin Avoid contact with eves skin or clothing Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse.

FIRST AID: If On Skin or Clothing

•Take off contaminated clothing.

•Rinse skin immediately with plenty of water for 15-20 minutes

•Call a poison control center or doctor for treatment advice

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic organisms, including fish and invertebrates. Do not apply directly to or near water. Drift and runoff may be hazardous to fish in water adjacent to treated areas. Do not contaminate water when disposing of equipment, washwaster, or rinsate. See Directions for Use for additional precautions and requirements. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging the treatment area.

PHYSICAL HAZARDS:

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130° F may cause bursting.

Contains No CFC's or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.

DIRECTIONS FOR USE:

To the extent consistent with applicable law. Seller makes no warranty, expressed or implied, concerning the use and handling of this product other than indicated on the label. To the extent consistent with applicable law. Buyer assumes all risks of use and handling of this material when such use and handling are contrary to label instructions.

USE ON	TO CONTROL
Tomatoes	Flea Beetles, Hornworms, Colorado Potato Beetles, Stink Bugs,
	Green Peach Aphids, Two Spotted Spider Mites, Whiteflies, Thrips
Peppers,	Colorado Potato Beetles, Leafhoppers, Blister Beetles,
Eggplants,	Green Peach Aphids, Flea Beetles, Thrips
Potatoes	
Cucumbers,	Striped and Spotted Cucumber Beetles, Aphids,
Squash, Melons	Two Spotted Spider Mites, Thrips
Beans	Mexican Bean Beetles, Whiteflies,
	12-Spotted Cucumber Beetles, Two Spotted Spider Mites, Thrips
Sweet Corn	Japanese Beetles
Carrots	Leafhoppers, Thrips
Peas	Aphids, Thrips
Cabbage	Aphids, Cabbage Loopers, Thrips

STORAGE AND DISPOSAL:

PESTICIDE STORAGE: Store in a cool area away from heat or open flame PESTICIDE DISPOSAL AND CONTAINER HANDLING: Do Not Puncture or Incinerate! IF EMPTY: Place in trash or offer for recycling if available. IF PARTLY FILLED: Call your local solid waste agency for disposal instructions.

Manufactured for: Ultrasol Industries Ltd. 10755 - 69 Avenue, Edmonton Alberta Canada T6H 2C9 780-432-6535



It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Read entire label and apply as directed. SHAKE WELL BEFORE USING.

USE RESTRICTIONS:

Do not allow adults, children, or pets to enter the treated area until sprays have dried. Do not make applications during rain. Do not wet plants to point of runoff or drip. Do not water the treated area to the point of runoff. All outdoor applications must be limited to spot treatments only, except for the following permitted

11565 1) Treatment to soil or vegetation around structures: 2) Applications to lawns, turf, and other vegetation To apply at the maximum rate, spray for 6 seconds to cover a space of 54 cubic feet (3' x 3' x 6' high). For best results, use throughout the growing season to prevent buildup of insect populations.

Sprav plants from a distance of 31/2 to 4 feet using short bursts (one second). To prevent injury to plants, avoid visibly wetting the foliage.

Do not apply this product in a way that will contact adults, children, or pets, either directly or through drift. Apply in the early morning or late evening when the air is still, treating all foliage with particular attention to the undersides of the leaves. Contact insects directly whenever possible. Repeat applications at 7 to 10 day intervals or after

heavy rains. Do not apply to food crops within one day of harvest.